

# Categorifying orthosymplectic Kazhdan–Lusztig polynomials

Ben Mills

University of York

30 Oct 2025


# A parable



# An alternative parable




## Silly shephards



- $|V \amalg W| = |V| + |W|$   
 $|V \times W| = |V| \cdot |W|$

# Jones polynomial


- In the 1980's Jones created a polynomial that was invariant under Reidemeister moves.



- It is not a complete invariant and it cannot detect the unknot.
- Khovanov 'categorified' the Jones polynomial by creating a cochain of graded vector spaces,  $H_n^n$ .
- The Jones polynomial appears in the cohomology of this cochain.
- This can detect the unknot.


# Generalised Khovanov arc algebras

- Brudan-Stroppel generalised this algebra,  $H_n^m$ , and associated it and its quasi-hereditary cover to blocks of parabolic category  $\mathcal{O}(m, n)$ .
- Related to the category of perverse sheaves on the Grassmannian of  $m$ -dimensional subspaces in an  $(m + n)$ -dimensional complex vector space.



# Generalised Khovanov arc algebras

- Brudan-Stroppel generalised this algebra,  $H_n^m$ , and associated it and its quasi-hereditary cover to blocks of parabolic category  $\mathcal{O}(m, n)$ .
- Related to the category of perverse sheaves on the Grassmannian of  $m$ -dimensional subspaces in an  $(m + n)$ -dimensional complex vector space.



- Ehrig-Stroppel generalised this generalised algebra into type  $D$  as well.

# Diagrammatic Hecke category

# Kazhdan-Lusztig polynomials

- In 1979, Kazhdan and Lusztig introduced the Kazhdan-Lusztig basis of the Hecke algebra of a Coxeter system. The base change matrix between the standard and the Kazhdan-Lusztig basis to be given by a set of polynomials  $n_{\mu,\lambda}(q)$ .

# Kazhdan-Lusztig polynomials

- In 1979, Kazhdan and Lusztig introduced the Kazhdan-Lusztig basis of the Hecke algebra of a Coxeter system. The base change matrix between the standard and the Kazhdan-Lusztig basis to be given by a set of polynomials  $n_{\mu,\lambda}(q)$ .
- There are many places these polynomials appear all over Representation Theory.

# Kahzdan-Lustzig polynomials

- In 1987, Deodhar introduced parabolic Kazhdan-Lusztig polynomials. These polynomials are defined starting from the choice of a Coxeter group, a standard parabolic subgroup and a sign.

# Kahzdan-Lustzig polynomials

- In 1987, Deodhar introduced parabolic Kazhdan-Lusztig polynomials. These polynomials are defined starting from the choice of a Coxeter group, a standard parabolic subgroup and a sign.
- In 1990, Deodhar proposed a (non-recursive) combinatorial approach to studying these polynomials.

# Kazhdan-Lusztig polynomials

- In 1987, Deodhar introduced parabolic Kazhdan-Lusztig polynomials. These polynomials are defined starting from the choice of a Coxeter group, a standard parabolic subgroup and a sign.
- In 1990, Deodhar proposed a (non-recursive) combinatorial approach to studying these polynomials.
- Elias-Williamson and Libedinsky-Williamson categorified the anti-spherical Kazhdan-Lusztig polynomials by interpreting them as composition factor multiplicities of simple modules within standard modules for the anti-spherical diagrammatic Hecke category,  $\mathcal{H}_{(W,P)}$  over  $\mathbb{C}$ .

# Kazhdan-Lusztig polynomials


- In 1987, Deodhar introduced parabolic Kazhdan-Lusztig polynomials. These polynomials are defined starting from the choice of a Coxeter group, a standard parabolic subgroup and a sign.
- In 1990, Deodhar proposed a (non-recursive) combinatorial approach to studying these polynomials.
- Elias-Williamson and Libedinsky-Williamson categorified the anti-spherical Kazhdan-Lusztig polynomials by interpreting them as composition factor multiplicities of simple modules within standard modules for the anti-spherical diagrammatic Hecke category,  $\mathcal{H}_{(W,P)}$  over  $\mathbb{C}$ .
- This proved the famous Kazhdan-Lusztig positivity conjecture

# Kazhdan-Lusztig polynomials

- In 1987, Deodhar introduced parabolic Kazhdan-Lusztig polynomials. These polynomials are defined starting from the choice of a Coxeter group, a standard parabolic subgroup and a sign.
- In 1990, Deodhar proposed a (non-recursive) combinatorial approach to studying these polynomials.
- Elias-Williamson and Libedinsky-Williamson categorified the anti-spherical Kazhdan-Lusztig polynomials by interpreting them as composition factor multiplicities of simple modules within standard modules for the anti-spherical diagrammatic Hecke category,  $\mathcal{H}_{(W,P)}$  over  $\mathbb{C}$ .
- This proved the famous Kazhdan-Lusztig positivity conjecture
- Williamson also used this structure to find an explicit counter-example to Lusztig's conjecture over  $\mathbb{F}_p$ .

## Maximal parabolic of type $(D_n, A_{n-1})$


- Take the Weyl group  $A_{n-1}$ , the subgroup of  $D_n (\leq C_n)$  generated by the reflections  $\{s_1, s_2, \dots, s_{n-1}\}$ .



- We can visualise the cosets  ${}^P W$  of this subgroup via their action permuting  $n$  labels from  $\{\wedge, \vee\}$ .
- In particular  $s_i$  acts by permuting the labels in position  $i$  and  $i + 1$  for  $1 \leq i \leq n - 1$ .
- $s_0$  permutes and flips (through the horizontal line) the labels in the first and second positions

## Example

- For example for  $n=4$ ,  $s_2$  acts as




## Example

- For example for  $n=4$ ,  $s_2$  acts as




## Example

- But  $s_0$  acts like:




## Example


- But  $s_0$  acts like:




# Labelling of cosets




# Labelling of cosets



# Labelling of cosets



# Labelling of cosets



# The diagrammatic Hecke category

## Definition

Define the Soergel generators to be the framed graphs



for reflections  $\sigma, \tau \in S_w$  for  $m(\sigma, \tau) = 2$ . Diagrams made by concatenation are called Soergel diagrams.

The diagrams represent morphisms between Soergel bimodules.

# The diagrammatic Hecke category

## Definition

Define the Soergel generators to be the framed graphs



for reflections  $\sigma, \tau \in S_w$  for  $m(\sigma, \tau) = 2$ . Diagrams made by concatenation are called Soergel diagrams.

The diagrams represent morphisms between Soergel bimodules.

## Definition

The Hecke category,  $\mathcal{H}_{(W,P)}$ , is spanned by all Soergel diagrams generated by the graphs above modulo a (medium) set of relations. The relations come from the Coxeter-Dynkin diagram of  $(W, P)$ .

# What can we say about the structure of $\mathcal{H}_{(W,P)}$ ?

- Labels of the highest weight modules:

$$\{\Delta(\mu) \mid \mu \in {}^P W\}$$

## What can we say about the structure of $\mathcal{H}_{(W,P)}$ ?

- Labels of the highest weight modules:

$$\{\Delta(\mu) \mid \mu \in {}^P W\}$$

- Labels of the simple modules:

$$\{L(\lambda) \mid \lambda \in {}^P W, \}$$

# What can we say about the structure of $\mathcal{H}_{(W,P)}$ ?

- Labels of the highest weight modules:

$$\{\Delta(\mu) \mid \mu \in {}^P W\}$$

- Labels of the simple modules:

$$\{L(\lambda) \mid \lambda \in {}^P W, \}$$

- Decomposition matrix:

$$n_{\mu,\lambda}(q) = \sum_{k \in \mathbb{Z}} [\Delta(\mu) : L(\lambda)\langle k \rangle] q^k$$

is uni-triangular wrt the partial ordering of  $\mathcal{G}_{(W,P)}$ .

# Calculating Type $D$ Kazhdan–Lusztig polynomials

# Cup diagrams

- This origin of this combinatorics starts with Lascoux-Schutzenberger and more concretely with Cox-De Vishcer and Lejczyk-Stroppel.

# Cup diagrams

- This origin of this combinatorics starts with Lascoux-Schutzenberger and more concretely with Cox-De Vishcer and Lejczyk-Stroppel.
- To a weight  $\mu \in {}^P W$  we create a cup diagram by drawing cups and strands underneath  $\mu$ .




# Oriented cup diagrams

- A cup diagram is oriented if:
  - For all undecorated cups one label points into the strand and one points out.
  - For all decorated cup connect nodes either both labels point in or both labels point out of the strand.
  - Every decorated cup or strand must be able to be deformed to the left edge.
  - Any strand from a  $\wedge$  node must be decorated.
  - Any strand from a  $\vee$  node must not.





# Oriented cup diagrams

- A cup diagram is oriented if:
  - For all undecorated cups one label points into the strand and one points out.
  - For all decorated cup connect nodes either both labels point in or both labels point out of the strand.
  - Every decorated cup or strand must be able to be deformed to the left edge.
  - Any strand from a  $\wedge$  node must be decorated.
  - Any strand from a  $\vee$  node must not.
- For example:







# Oriented cup diagrams

- A cup diagram is oriented if:
  - For all undecorated cups one label points into the strand and one points out.
  - For all decorated cup connect nodes either both labels point in or both labels point out of the strand.
  - Every decorated cup or strand must be able to be deformed to the left edge.
  - Any strand from a  $\wedge$  node must be decorated.
  - Any strand from a  $\vee$  node must not.
- For example:






# Oriented cup diagrams

- A cup diagram is oriented if:
  - For all undecorated cups one label points into the strand and one points out.
  - For all decorated cup connect nodes either both labels point in or both labels point out of the strand.
  - Every decorated cup or strand must be able to be deformed to the left edge.
  - Any strand from a  $\wedge$  node must be decorated.
  - Any strand from a  $\vee$  node must not.
- For example:






# Oriented cup diagrams

- A cup diagram is oriented if:
  - For all undecorated cups one label points into the strand and one points out.
  - For all decorated cup connect nodes either both labels point in or both labels point out of the strand.
  - Every decorated cup or strand must be able to be deformed to the left edge.
  - Any strand from a  $\wedge$  node must be decorated.
  - Any strand from a  $\vee$  node must not.
- For example:



# Oriented cup diagrams

- A cup diagram is oriented if:
  - For all undecorated cups one label points into the strand and one points out.
  - For all decorated cup connect nodes either both labels point in or both labels point out of the strand.
  - Every decorated cup or strand must be able to be deformed to the left edge.
  - Any strand from a  $\wedge$  node must be decorated.
  - Any strand from a  $\vee$  node must not.
- For example:



# Oriented cup diagrams

- Define the degree of a cup diagram  $D$  to be:


$$\# \left\{ \begin{array}{c} \curvearrowright \\ \cup \end{array} , \begin{array}{c} \cup \\ \bullet \\ \cup \end{array} \right\}$$

# Oriented cup diagrams

- Define the degree of a cup diagram  $D$  to be:

$$\# \left\{ \begin{array}{c} \curvearrowright \\ \cup \\ \curvearrowright \end{array}, \begin{array}{c} \cup \\ \cup \\ \bullet \end{array} \right\}$$

- For example:





## Neutral elements

- For  $\mu \in {}^P W$  we want to construct the unique maximal cup  $\underline{\mu}$  of degree zero.

$$\# \left\{ \begin{array}{c} \text{Y} \\ \text{Y} \end{array} \cup, \begin{array}{c} \text{Y} \cup \\ \text{Y} \end{array} \right\}$$

- There is a nice algorithm for us:




## Neutral elements

- For  $\mu \in {}^P W$  we want to construct the unique maximal cup  $\underline{\mu}$  of degree zero.

$$\# \left\{ \begin{array}{c} \curvearrowright \\ \cup \end{array}, \begin{array}{c} \cup \\ \curvearrowleft \end{array} \right\}$$

- There is a nice algorithm for us:




## Neutral elements

- For  $\mu \in {}^P W$  we want to construct the unique maximal cup  $\underline{\mu}$  of degree zero.

$$\# \left\{ \begin{array}{c} \text{Y} \\ \cup \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \cup \\ \bullet \\ \cup \\ \text{Y} \end{array} \right\}$$

- There is a nice algorithm for us:




# Neutral elements

- For  $\mu \in {}^P W$  we want to construct the unique maximal cup  $\underline{\mu}$  of degree zero.

$$\# \left\{ \begin{array}{c} \text{Y} \\ \cup \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \text{ Y} \\ \cup \\ \bullet \end{array} \right\}$$

- There is a nice algorithm for us:




## Neutral elements

- For  $\mu \in {}^P W$  we want to construct the unique maximal cup  $\underline{\mu}$  of degree zero.

$$\# \left\{ \begin{array}{c} \curvearrowright \\ \cup \end{array}, \begin{array}{c} \cup \\ \curvearrowleft \end{array} \right\}$$

- There is a nice algorithm for us:







## Neutral elements









- For  $\mu \in {}^P W$  we want to construct the unique maximal cup  $\underline{\mu}$  of degree zero.










$$\# \left\{ \begin{array}{c} \curvearrowright \\ \cup \end{array}, \begin{array}{c} \cup \\ \curvearrowright \end{array} \right\}$$
















- There is a nice algorithm for us:




















































.








.










.





.



.








.




.





.

.









.


.

























.



.



								
		.	.	.	.	.	.	.
			.	.	.	.	.	.
	.			.	.	.	.	.
	.	.			.	.	.	.
	.	.		.		.	.	.
	.						.	.
			.	.	.			.
		.	.	.	.	.		



.

.



.

.

.

.

.



.


.

.



.

.

.



.



.

.

.



.

.



.

.



.

.


.

.




.

.



.






.

.

.




.



.


.




.

.

.



.





.























.





.

.

.




								
		.	.	.	.	.	.	.
			.	.	.	.	.	.
	.			.	.	.	.	.
	.	.			.	.	.	.
	.	.		.		.	.	.
	.						.	.
			.	.	.			.
		.	.	.	.	.		

								
	1	.	.	.	.	.	.	.
	$q$	1	.	.	.	.	.	.
	.	$q$	1	.	.	.	.	.
	.	.	$q$	1	.	.	.	.
	.	.	$q$	.	1	.	.	.
	.	$q$	$q^2$	$q$	$q$	1	.	.
	$q$	$q^2$	.	.	.	$q$	1	.
	$q^2$	.	.	.	.	.	$q$	1

# Flip it

- Every oriented cup diagram of degree  $k$  is created by ‘flipping’  $k$  cups in the diagram  $\underline{\mu\mu}$


$$\deg(D) = \# \left\{ \begin{array}{c} \curvearrowright \\ \cup \\ \curvearrowleft \end{array} + \begin{array}{c} \curvearrowleft \\ \cup \\ \curvearrowright \end{array} \right\}$$



# Flip it

- Every oriented cup diagram of degree  $k$  is created by ‘flipping’  $k$  cups in the diagram  $\underline{\mu\mu}$


$$\text{deg}(D) = \# \left\{ \begin{array}{c} \curvearrowright \\ \cup \\ \curvearrowleft \end{array} + \begin{array}{c} \curvearrowleft \\ \cup \\ \curvearrowright \end{array} \right\}$$



# Flip it

- Every oriented cup diagram of degree  $k$  is created by ‘flipping’  $k$  cups in the diagram  $\underline{\mu\mu}$


$$\text{deg}(D) = \# \left\{ \begin{array}{c} \curvearrowright \\ \cup \\ \curvearrowleft \end{array} + \begin{array}{c} \cup \\ \curvearrowright \\ \bullet \\ \cup \\ \curvearrowleft \end{array} \right\}$$



# Flip it

- Every oriented cup diagram of degree  $k$  is created by ‘flipping’  $k$  cups in the diagram  $\underline{\mu\mu}$

$$\deg(D) = \# \left\{ \begin{array}{c} \curvearrowright \\ \cup \\ \bullet \end{array} + \begin{array}{c} \cup \\ \curvearrowright \\ \bullet \end{array} \right\}$$



# Generators

- So calculating Kazhdan–Lusztig polynomials (or equivalently composition factors of highest weight modules in the Hecke category) is really only the simple matter of flipping cups in the diagram  $\underline{\mu\mu}$ .

# Generators

- So calculating Kazhdan–Lusztig polynomials (or equivalently composition factors of highest weight modules in the Hecke category) is really only the simple matter of flipping cups in the diagram  $\underline{\mu\mu}$ .
- It would be ‘nice’ if this cup flipping could control even more....

# Generators

## Definition

We say  $\lambda = \mu - p$  when  $\underline{\lambda}\lambda$  and  $\underline{\mu}\mu$  differ by the flipping of a single cup  $p$ .

## Theorem (M. 25+)

*The algebra  $\mathcal{H}_{(D_n, A_{n-1})}$  is the associative  $\mathbb{k}$ -algebra generated by the elements*

$$\{\underline{\mu}\lambda \mid \lambda, \mu \in {}^P W \text{ with } \lambda = \mu - p \text{ for some } p \in \underline{\mu}\} \cup \{\underline{\mu}\mu \mid \mu \in {}^P W\}$$

*modulo (a medium set of) degree two relations.*

# Generators

## Definition

We say  $\lambda = \mu - p$  when  $\underline{\lambda}\lambda$  and  $\underline{\mu}\mu$  differ by the flipping of a single cup  $p$ .

## Theorem (M. 25+)

*The algebra  $\mathcal{H}_{(D_n, A_{n-1})}$  is the associative  $\mathbb{k}$ -algebra generated by the elements*

$$\{\underline{\mu}\lambda \mid \lambda, \mu \in {}^P W \text{ with } \lambda = \mu - p \text{ for some } p \in \underline{\mu}\} \cup \{\underline{\mu}\mu \mid \mu \in {}^P W\}$$

*modulo (a medium set of) degree two relations.*

- This gives us a purely combinatorial presentation of  $\mathcal{H}_{(D_n, A_{n-1})}$  that is positively graded, quadratic, modulo relations that are defined entirely on the cup diagrams.

# Generators

## Definition

We say  $\lambda = \mu - p$  when  $\underline{\lambda}\lambda$  and  $\underline{\mu}\mu$  differ by the flipping of a single cup  $p$ .

## Theorem (M. 25+)

*The algebra  $\mathcal{H}_{(D_n, A_{n-1})}$  is the associative  $\mathbb{k}$ -algebra generated by the elements*

$$\{\underline{\mu}\lambda \mid \lambda, \mu \in {}^P W \text{ with } \lambda = \mu - p \text{ for some } p \in \underline{\mu}\} \cup \{\underline{\mu}\mu \mid \mu \in {}^P W\}$$

*modulo (a medium set of) degree two relations.*

- This gives us a purely combinatorial presentation of  $\mathcal{H}_{(D_n, A_{n-1})}$  that is positively graded, quadratic, modulo relations that are defined entirely on the cup diagrams. Over a field, this gives an Ext-quiver presentation.

# Generators

## Definition

We say  $\lambda = \mu - p$  when  $\underline{\lambda}\lambda$  and  $\underline{\mu}\mu$  differ by the flipping of a single cup  $p$ .

## Theorem (M. 25+)

*The algebra  $\mathcal{H}_{(D_n, A_{n-1})}$  is the associative  $\mathbb{k}$ -algebra generated by the elements*

$$\{\underline{\mu}\lambda \mid \lambda, \mu \in {}^P W \text{ with } \lambda = \mu - p \text{ for some } p \in \underline{\mu}\} \cup \{\underline{\mu}\mu \mid \mu \in {}^P W\}$$


*modulo (a medium set of) degree two relations.*

- This gives us a purely combinatorial presentation of  $\mathcal{H}_{(D_n, A_{n-1})}$  that is positively graded, quadratic, modulo relations that are defined entirely on the cup diagrams. Over a field, this gives an Ext-quiver presentation.
- This presentation allows us to define an explicit isomorphism into the generalised Khovanov arc algebra of type  $D$ ,  $\mathbb{D}_n$ .


We hope to utilise this to provide a faithful quasi-hereditary cover for  $\mathbb{D}_n$ .



Hecke category




Cup combinatorics




Generalized Khovanov arc diagrams



Hecke category



Cup combinatorics



Generalized Khovanov arc diagrams



## Structure of highest weight modules

- This also presentation tells us another way to compute the full submodule lattice of any highest weight module too
- There are nice properties for the structure of  $\Delta(\mu)$ :

# Structure of highest weight modules

- This also presentation tells us another way to compute the full submodule lattice of any highest weight module too
- There are nice properties for the structure of  $\Delta(\mu)$ :
  - The radical filtration coincides with the grading layers.

# Structure of highest weight modules

- This also presentation tells us another way to compute the full submodule lattice of any highest weight module too
- There are nice properties for the structure of  $\Delta(\mu)$ :
  - The radical filtration coincides with the grading layers.
  - One dimensional simple socle  $L(\mu_n)$ , indexed by a unique element of maximal degree  $\mu_n$ .

# Structure of highest weight modules

- This also presentation tells us another way to compute the full submodule lattice of any highest weight module too
- There are nice properties for the structure of  $\Delta(\mu)$ :
  - The radical filtration coincides with the grading layers.
  - One dimensional simple socle  $L(\mu_n)$ , indexed by a unique element of maximal degree  $\mu_n$ .
  - One dimension simple head  $L(\mu)$ .

# Structure of highest weight modules

- This also presentation tells us another way to compute the full submodule lattice of any highest weight module too
- There are nice properties for the structure of  $\Delta(\mu)$ :
  - The radical filtration coincides with the grading layers.
  - One dimensional simple socle  $L(\mu_n)$ , indexed by a unique element of maximal degree  $\mu_n$ .
  - One dimension simple head  $L(\mu)$ .
  - An extension only when you can ‘flip’ one cup to get between two cups diagrams:

We have edges

$$L_n(\mu) \longrightarrow L_n(\nu)$$

in an Alperin diagram whenever  $\mu = \nu \pm p$  for some  $p \in \underline{\nu}$  or  $p \in \underline{\mu}$ .

- Reminder: the composition factors  $[\Delta(\mu) : L(\mu)]\langle k \rangle$  give us the Kazhdan-Lusztig polynomials.

## Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$

# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$



# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$




# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$




# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$




# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$



# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$



# Structure of highest weight module

$$\Delta(s_0 s_2 s_3 s_4 s_5 s_1) \in \mathcal{H}_{(D_7, A_6)}$$

